

Ant-Tended Hemiptera in Amazonian Myrmecophytes:
Patterns of Abundance and Implications for Mutualism
Function (Hymenoptera: Formicidae)

433

by

David M. Lapola^{1,e}, Emilio M. Bruna^{2,3,a}, Cristina Granara de
Willink⁴, & Heraldo L. Vasconcelos^{5,f}

ABSTRACT

We assessed how the abundance of ant-tended Hemiptera associated with two Amazonian myrmecophytes, *Tococa bullifera* and *Mateia guianensis*, varied as a function of resident ant species. We collected five species or morpho-species of adult hemiptera in the domatia of *M. guianensis*, with four of these species also found in *Tococa bullifera*. *Mateia guianensis* plants inhabited by *Crematogaster laevis* had over four-fold more hemiptera in them than plants inhabited by *Pheidole murutuba*. In contrast, the density of hemiptera in *Tococa bullifera* domatia was independent of the species of ant resident. For each of the two ant species inhabiting *Mateia guianensis*, there was a positive and significant relationship between the abundance of hemiptera and workers inhabiting a plant. This relationship was also significant and positive for the *Tococa bullifera* plants inhabited by *C. laevis*. However, there was no relationship between *Azteca* worker and hemipteran density, although there was a trend towards a positive relationship. Our results indicate that hemipteran abundance can vary significantly between different myrmecophyte species, but that the nature of this relationship is mediated by the identity of the ant associate. Because hemipterans are herbivores, the costs and benefits of different ant partners to the host plant may vary in ways that are often overlooked.

¹Departamento de Ecologia, Universidade Estadual Paulista, Rio Claro, SP, 13506-900 Brazil

²Department of Wildlife Ecology and Conservation, University of Florida, Gainesville, FL, 32611-0430 USA

³Tropical Conservation and Development Program, Center for Latin American Studies, University of Florida, Gainesville, FL, 32611-0430 USA

⁴Instituto Superior de Entomología, Universidad Nacional de Tucumán, Miguel Lillo 205, 4000 San Miguel de Tucumán, Argentina

⁵Instituto de Biología, Universidade Federal de Uberlândia, Uberlândia, MG, 38400-902 Brazil
⁶Biological Dynamics of Forest Fragments Project, INPA-PDBFF, CP 478, 69011-970 Manaus, AM, Brazil

^a Author and address for correspondence: Department of Wildlife Ecology and Conservation, University of Florida, PO Box 110430, Gainesville, FL 32611-0430, USA. Phone: (352) 946-0634, Fax: (352) 392-6994, Email: BrunaE@wec.ufl.edu

434

Sociobiology Vol. 46, No. 2, 2005

Keywords: ant-plants, *Azteca* coccids, *Crematogaster laevis*, *Mateia guianensis*, mealybugs, Melastomataceae, pseudococcids, scale insects, *Tococa bullifera*.

INTRODUCTION

Mutualistic associations between plants and ants have long been recognized as one of the defining features of tropical forests (Beattie 1985, Huxley & Cutler 1991). In some cases the mutualisms are facultative – ants may simply forage on leaf surfaces or feed from extra-floral nectaries, with varying degrees of specialization. However, many ant species obligately establish colonies in hollow thorns, swollen petioles, leaf pouches, or other specialized plant structures known as domatia. Because ant residents may confer benefits such as defense against herbivores, they have been the subject of considerable observational and experimental research (reviewed in Beattie 1985, Huxley & Cutler 1991, Bronstein 1998, Heil & Mckey 2003).

Obligate ant-plant mutualisms almost always involve a third partner: scale insects and mealybugs (coccids and pseudococcids, respectively, both Hemiptera). These herbivores are often tended by ants, which use the "honeydew" they excrete as a food resource (Way 1963, Carroll & Janzen 1973). Their presence putatively has both costs and benefits to the host plant – they feed on sap and are potential vectors for plant diseases (McPherson & Chung-Kim 1993), but they can also influence the number of ants patrolling leaves via their influence on ant behavior and colony structure (Buckley 1987). They can therefore alter the outcome of plant-ant interactions along the continuum from mutualism to parasitism (Gaume *et al.* 1998).

There is an extensive body of literature on ant-plant-hemipteran interactions (reviewed in Way 1963, Buckley 1987), motivated in part by their importance as agricultural pests. However, few of these studies experimentally investigate the ecology of hemiptera associated with obligate domatia-dwelling ants (but see Gaume *et al.* 1998, Heckroth *et al.* 1998, Gaume *et al.* 2000, Itino *et al.* 2001, Gaume & Mckey 2002). This is due primarily to the difficulty of conducting manipulations with domatia residents and the need for destructive sampling of ant colonies. An additional complicating factor is that we have only a limited understanding of the taxonomy, systematics, distribution, and abundance of most hemiptera found in ant-plants. Such data are an important prerequisite for investigating how hemiptera may influence the dynamics of ant-plant mutualisms.

In this study we investigated the relationship between the ants and hemiptera associated with two Amazonian myrmecophytes – *Mateia*

guyanensis and *Tococa bullifera* (both Melastomataceae). As with most ant-plant species, these species are associated with a suite of ant species (Fowler 1993, Fonseca & Ganade 1996) though each individual plant is colonized by a single species at a time (Vasconcelos and Davidson 2000). Our previous work in these systems has demonstrated important interspecific variation in several aspects of plant-ant interactions (Lapola et al. 2003, Bruna et al. 2004), and we expect the same is probably true of the relationship between ants, plants, and hemiptera as well. With this in mind, we addressed the following three questions: First, what hemiptera species are associated with *T. bullifera* and *M. guianensis*? Second, does hemipteran abundance vary with respect to the identity of host-plants and their ant residents? Third, does the abundance of hemiptera vary with respect to the ant-worker abundance?

MATERIALS AND METHODS

Study site and system

Fieldwork was conducted from January to February 2004 in Reserve 1501 of the Biological Dynamics of Forest Fragments Project (BDFFP), located c. 70 km north of Manaus (2° 25'S, 59° 48'W). This reserve of approximately 800 ha is embedded in more than 10,000 ha of continuous terra-firme rain forest. Annual precipitation ranges from 1900–3500 mm, with a distinct dry season from May–December. For a complete description of the site see Bierregaard et al. (2002).

Maieta guianensis and *Tococa bullifera* are two of the sixteen species of myrmecophytes that have been identified in Reserve 1501 (Fonseca & Ganade 1996, Bruna et al. 2005). Both are understory shrubs, reaching heights of 2 and 3 m, respectively, with swollen pouches at the bases of leaves in which ant queens establish colonies. In our study sites, approximately 30% of the *T. bullifera* individuals and 14% of the *M. guianensis* individuals are occupied by *Crematogaster laevis*. The remaining 86% of *M. guianensis* and 70% of *Tococa bullifera* are occupied by *Pheidole minutula* and an unidentified species of *Azteca*, respectively (Vasconcelos & Davidson 2000).

Data collection and statistical analysis

To elucidate the relationship between plants, ants, and hemiptera, we collected five randomly selected domatia from 10 *T. bullifera* and 10 *M. guianensis* inhabited by each ant species (total $N = 40$ plants). For each plant we counted (a) the total number of ant workers in the five domatia and (b) the total number of number of adult hemiptera. We identified hemiptera to species or morpho-species: voucher specimens are deposited in the entomological collection of Brazil's Instituto Nacional de Pesquisas da Amazônia (INPA).

To compare the abundance of (a) hemiptera in plants inhabited by each ant species and (b) ant workers of different species inhabiting the same plant species, we summed the number of hemiptera and ant workers in the five domatia collected from each plant and compared the total abundance using *t*-tests. We log-transformed all totals prior to analysis to meet the assumptions of parametric statistics, but throughout the manuscript we present back-transformed values. Each plant species was compared separately. To test for a relationship between hemipteran abundance and worker abundance (both log-transformed) we used simple linear regression. Each plant versus ant species combination was compared separately.

RESULTS

We collected five species or morpho-species of adult hemiptera in the domatia of *M. guianensis*: *Nipaeococcus* morpho-species 1, *Nipaeococcus* morpho-species 2, *Dysmicoccus* morpho-species 1 (all Pseudococcidae), *Coccus viridis*, and *Coccus hesperidum* (both Coccidae). Although the identification of immature individuals is difficult, we also collected immature individuals that we are provisionally assigning to the genus *Alchertesia* and the family Margarodidae. All of these species were also found in the domatia of *T. bullifera*, with the exception of *Nipaeococcus* morpho-species 1.

Maieta guianensis plants inhabited by *Crematogaster laevis* had over four-fold more hemiptera in them than plants inhabited by *Pheidole minutula* (231.6 ± 88.07 SE vs. 49.1 ± 6.63 SE, respectively, Fig. 1A), a highly significant difference ($t=3.301$, $df = 18$, $P = 0.004$). In contrast, the density of hemiptera in *Tococa bullifera* domatia was independent of the species of ant resident ($t=0.498$, $df = 18$, $P = 0.62$). The average abundance of hemipterans in plants inhabited by *C. laevis* was 82.1 ± 22.22 SE, while in plants inhabited by *Azteca* sp. it was 94.5 ± 26.78 SE (Fig. 1A).

The summed abundance of *C. laevis* workers collected from the domatia of *T. bullifera* plants was over 1.5 times that of *Azteca* sp. workers (295.8 ± 41.48 SE vs. 186.6 ± 52.43 SE, respectively, Fig. 1B). A similar pattern was seen in *Maieta guianensis*—plants with *C. laevis* colonies in them had twice the number of workers in domatia as those inhabited by *P. minutula* (208.5 ± 28.33 SE vs. 130.8 ± 16.28 SE, respectively). However, the difference in worker abundance was only significant for *T. bullifera* (*T. bullifera*: $t = 2.27$, $df = 18$, $P = 0.04$; *M. guianensis*: $t = 0.63$, $df = 18$, $P = 0.09$, Fig. 1B).

For each of the two ant species inhabiting *Maieta guianensis*, there was a positive and significant relationship between the abundance of hemiptera and workers inhabiting a plant (*C. laevis*: $MS = 0.849$, $F_{1,8} =$




Fig. 1. Summed abundance of (A) hemiptera and (B) ant workers in five domatia inhabited by different ant species colonizing the Amazonian myrmecophytes *Maletia guianensis* and *Tococa bullifera*. Comparisons for each plant species were conducted separately (see text for details).

6.07, $P = 0.04$, $R^2 = 0.43$; $P. minutula$: $MS=0.849$, $F_{1,8} = 6.07$, $P = 0.04$, $R^2 = 0.43$, Fig. 2A). This relationships was also significant and positive for the *Tococa bullifera* plants inhabited by *C. laevis* ($MS=1.88$, $F_{1,8} = 6.69$, $P = 0.04$, $R^2 = 0.46$, Fig. 2B). However, there was no relationship between *Azteca* worker and hemipteran density, although there was a trend towards a positive relationship ($MS=1.56$, $F_{1,8} = 2.40$, $P = 0.16$, $R^2 = 0.23$).

DISCUSSION

Domatia-dwelling hemiptera are a common partner in obligate ant-plant mutualisms. The ecology of these interactions continues to be understudied, however, particularly in the Neotropics. While studies of *Maccaranga* trees in southeast Asia have suggested that some hemipteran species may have obligate associations with single plant species (Heckroth et al. 1998), two of the species we collected – *Coccus hesperidum* (i.e., the soft brown scale) and *Coccus viridis* (i.e., the green soft scale) – have wide host ranges and cosmopolitan geographic distributions (Delte & Fasulo 2001). Much less is known regarding the host-plant preference and geographic range of the other genera we collected (e.g., *Nipaecoccus*, *Dysmicoccus*), despite their importance as agricultural pests (reviewed in Ben-Dov & Hodgson 1997). Comprehensive surveys of the hemiptera associated with Neotropical ant-plants, such as those Heckroth et al. (1998) have conducted with *Maccaranga*, are clearly needed.

Plant identity, ant identity, and the abundance of hemiptera

Our results indicate that hemipteran abundance can vary significantly between different myrmecophyte species, but that the nature of




Fig. 2. Relationship between hemipteran and ant worker abundance in (A) *Maletia guianensis* and (B) *Tococa bullifera* plants inhabited by different ant species. For *M. guianensis* the linear regression equations describing the relationships are: $\ln(\text{no. of } P. minutula \text{ workers}) = 2.62 + 0.57 * \ln(\text{no. of hemiptera})$, $R^2 = 0.44$; $\ln(\text{no. of } C. laevis \text{ workers}) = 3.72 + 0.31 * \ln(\text{no. of hemiptera})$, $R^2 = 0.43$. For *T. bullifera* the equations are: $\ln(\text{no. of } Azteca \text{ sp. workers}) = 2.67 + 0.52 * \ln(\text{no. of hemiptera})$, $R^2 = 0.23$; $\ln(\text{no. of } C. laevis \text{ workers}) = 3.49 + 0.50 * \ln(\text{no. of hemiptera})$, $R^2 = 0.46$.

this relationship is mediated by the identity of the ant associate. *Maletia guianensis* plants inhabited by *C. laevis* colonies had four times more hemiptera in them than those inhabited by *P. minutula*. This suggests *C. laevis* is more heavily dependent on hemipteran honeydew as a food resource than *P. minutula*, a conclusion that is the consistent with the results of previous work conducted in this system. Vasconcelos and Davidson (2000) found that the proportion of nitrogen in *C. laevis* workers was lower than in workers of *P. minutula*, indicating animal protein made up a lower proportion of their diet. Because as much as 25% of a *C. laevis* colony may be in satellite nests in twigs at the base of the plant (Vasconcelos, unpubl. data) some of the honeydew collected in domatia may also be going to workers without access to hemipterans.

Why then is the number of hemiptera tended by *C. laevis* so much lower when this species inhabits *T. bullifera*? At least three non-mutually exclusive mechanisms could explain these results. First, insect prey may be more abundant on the larger *T. bullifera* plants than on *M. guianensis* (e.g., *T. bullifera* may be more "apparent", sensu Feeny 1976) thereby reducing the dependence of *C. laevis* on hemipteran honeydew. Alternatively, the glandular trichomes on the interior surface of *T. bullifera*'s domatia may serve as an alternative source of carbohydrates (Roth 1976), thereby reducing the need to tend large numbers of hemiptera. Finally, we cannot discount the possibility that our results reflect the relatively limited number of colonies we sampled ($N = 10$ for each species x host plant combination). It may be that hemipteran abundance varies as a function of colony characteristics we did not control (e.g., colony age, microhabitat), and that increased

sampling would reduce the disparity between *C. laevis* colonies in different host-plant species. Regardless of the mechanism, our results suggest the relationship between domatia-inhabiting ants and hemiptera may be context-dependent in ways that have previously remained unexplored. While earlier comprehensive studies (e.g., Fonseca 1993, Gaume et al. 1998, Gaume et al. 2000, Gaume & Mckey 2002) have investigated the causes and consequences of variation in hemipteran abundance within a single plant species, most plant-ants are actually associated with a suite of geographically co-occurring taxa (e.g., Heckroth et al. 1998). Our results clearly indicate that conclusions from studies conducted with one plant species may not readily be generalized to others, even when these species are closely related.

Is ant colony size limited by hemipteran abundance?

The size of ant colonies in myrmecophytic plants are thought to be regulated by two principal factors: nesting space (i.e., domatia volume) or the availability of food (Hölldobler & Wilson 1990, Gaume et al. 1998). Previous studies investigating how honeydew, food bodies, or other food resources influence the size of ant colonies associated with myrmecophytes have drawn contradictory conclusions. While Fonseca (1993) found the volume of domatia most strongly limited colony size, other have found hemiptera appear to exert a strong influence on colony size and behavior (Gaume et al. 1998, Gaume & Mckey 2002). These patterns may in part be influenced by the source of nutritional resources – it has been argued that when the plant indirectly provides nutritional resources (i.e., via hemipteral), colonies are more likely to be limited by space rather than food (Gaume et al. 1998). The results of our analyses, however, are not consistent with this hypothesis. In three of four ant species x plant species combinations, there was a positive and significant relationship between ant worker abundance and hemipteran abundance. This suggests that in contrast to other myrmecophytic systems in our study sites (Fonseca 1993), the abundance of hemiptera and other food sources could potentially limit colony size. However, other factors are clearly important, since hemipteran abundance explained only 43-46% of the variance in worker number. We suggest future studies investigating the factors limiting plant-ant colony size move beyond the correlative framework used in this and other studies, and instead conduct experimental manipulations of food abundance and colony size.

Finally, interspecific variability in hemipteran tending has importance consequences for plants as well. Because mealybugs and scales are herbivores, the maintenance of high densities by ant colonies could

conceivably have costs for plant growth and reproduction. Our results suggest these costs would be similar for *T. bullifera* plants irrespective of their ant inhabitant, since both *C. laevis* and *Azeca* sp. tended similarly low numbers of hemiptera. However, the disparity in hemipteran abundance observed in *M. guianensis* suggests they may incur much higher costs when hosting *C. laevis* colonies than when inhabited by *P. nuttallia*. These costs include direct costs from hemipteran herbivory, as well as indirect costs resulting from reduced patrolling of leaves by ants (sensu Gaume et al. 1998). Such variable costs are frequently overlooked in studies evaluating the benefits to plants of their ant associates, and could conceivably diminish the purported benefits of ant colonies substantially.

CONCLUSION

Our study is one of a limited number to have compared the abundance of hemiptera found in the domatia of different myrmecophytic plant species (see also Heckroth et al. 1998, Itino et al. 2001). We found that the density of hemiptera varies within and between plant species, and that this can be a function of the identity of ant colonists. As such, the costs and benefits of different ant partners to the host plant may vary in ways that are often overlooked. This may help explain the observed disparities in colonization frequencies by the different ant species associated with ant-plants (Longino 1989, Fonseca & Ganade 1996, Bruna et al. 2005). Previous work has focused almost exclusively on ant defensive behavior, habitat specificity, and tradeoffs between competitive ability and queen dispersal as the principal drivers of distribution patterns (e.g., Yu & Davidson 1997, Stanton et al. 2002, Bruna et al. 2004). We propose the ubiquitous intermediate trophic level could be playing a critical role as well. We also suggest studying the ecology of coccids and pseudococcids may yield insights into the post-colonization dynamics of ant-plant mutualisms. The chemical composition of the honeydew ants consume can vary as a function of the location on the plant where insects are feeding, plant age, plant species, and season (reviewed in Way 1963, Molyneux et al. 1990). If tending, it may indirectly influence ant patrolling behavior and hence levels of herbivory. Finally, we end by encouraging other researchers working on tropical ant-plant mutualisms to at the very least collect voucher specimens of the coccids and pseudococcids inhabiting ant-plants. Information on the geographic distribution of these cryptic insects and their host-plant associations is a necessary first step in elucidating their impacts on the dynamics of ant-plant mutualisms.

ACKNOWLEDGMENTS

We would like to thank B. Ronchi-Teles for helpful discussions. The Biological Dynamics of Forest Fragments Project provided logistical support; financial support was provided by the National Science Foundation (Grants DEB-0309819 & DEB-0453631), the BDFFP, the University of Florida's Institute for Food and Agricultural Sciences, and the Brazilian Council of Research and Scientific Development (CNPq). This is publication number 441 in the BDFFP Technical Series and FAES Publication number R-10879.

LITERATURE CITED

- Beattie, A.J. 1985. The evolutionary ecology of ant-plant interactions. Cambridge University Press, Cambridge, UK.
- Ben-Dov, Y. & C.J. Hodgson (eds.) 1997. Scale insects: their biology, natural enemies and control. Elsevier, New York.
- Bierregaard, R.O., C. Gascon, T.E. Lovejoy, & R. Mesquita (eds.) 2002. Lessons from Amazonia: the ecology and conservation of a fragmented forest. Yale University Press, New Haven.
- Bronstein, J.L. 1998. The contribution of ant-plant protection studies to our understanding of mutualism. *Biotropica* 30: 150-161.
- Bruna, E.M., D.M. Lapola, & H.L. Vasconcelos 2004. Interspecific variation in the defensive responses of obligate plant-ants: experimental tests and consequences for herbivory. *Oecologia* 138: 558-565.
- Bruna, E.M., H.L. Vasconcelos, & S. Heredia 2005. The effect of habitat fragmentation on communities of mutualists: Amazonian ants and their host plants. *Biological Conservation* 124: 209-216.
- Buckley, R. 1987. Ant-plant-homopteran interactions. *Advances in Ecological Research* 16: 53-85.
- Carroll, C.R. & D.H. Janzen 1973. Ecology of foraging by ants. *Annual Review of Ecology and Systematics* 4: 231-257.
- DeJele, G.W. & T.R. Fasulo 2001. Green Scale, *Coccus viridis* (Green) (Insecta: Homoptera: Coccidae). Publication number FENY-253 of Florida Cooperative Extension Service, Institute for Agricultural Sciences, University of Florida.
- Feeny, P. 1976. Plant apparency and chemical defense. *Rec. Adv. Phytochem* 10: 1-40.
- Fonseca, C.R. 1993. Nesting space limits colony size of the plant-ant *Pseudomyrmex concolor*. *Oikos* 67: 473-482.
- Fonseca, C.R. & G. Ganade 1996. Asymmetries, compartments and null interactions in an Amazonian ant-plant community. *Journal of Animal Ecology* 65: 339-347.
- Fowler, H.G. 1993. Herbivory and assemblage structure of myrmecophytous understory plants and their associated ants in the central Amazon. *Insectes Sociaux* 40: 137-145.
- Gaume, L. & D. Mckey 2002. How identity of the homopteran trophobiont affects sex allocation in a symbiotic plant-ant: the proximate role of food. *Behavioral Ecology and Sociobiology* 51: 197-205.
- 442
- Sociobiology Vol. 46, No. 2, 2005
- Gaume, L., D. Mckey, & S. Terrin 1998. Ant-plant-homopteran mutualism: how the third partner affects the interaction between a plant-specialist ant and its myrmecophyte host. *Proceedings of the Royal Society of London Series B-Biological Sciences* 265: 569-575.
- Gaume, L., D. Mattie-Ferrero, & D. Mckey 2000. Colony formation and acquisition of coccoid trophobionts by *Aphoromyrmex qfer* (Formicidae): co-dispersal of queens and phoretic mealybugs in an ant-plant-homopteran mutualism? *Insectes Sociaux* 47: 84-91.
- Heckroth, H.P., B. Fiala, P.J. Gillian, A.H. Idhrs, & U. Maschwitz 1998. The soft scale (Coccidae) associates of Malaysian ant-plants. *Journal of Tropical Ecology* 14: 427-443.
- Hell, M., & D. Mckey 2003. Protective ant-plant interactions as model systems in ecological and evolutionary research. *Annual Review of Ecology, Evolution and Systematics* 34: 425-453.
- Hölldobler, B. & E.O. Wilson 1990. *The ants*. Belknap Press of Harvard University Press, Cambridge, MA.
- Huxley, C.R., & D.F. Cutler (eds.) 1991. *Ant-plant interactions*. Oxford University Press, Oxford.
- Ihno, T., T. Itohka, A. Hatada, & A.A. Hamid 2001. Effects of food rewards offered by ant-plant *Macaranga* on the colony size of ants. *Ecological Research* 16: 775-786.
- Lapola, D.M., E.M. Brunna, & H.L. Vasconcelos 2003. Contrasting responses to induction cues by ants inhabiting *Mateia guidonensis* (Melastomataceae). *Biotropica* 35: 295-300.
- Longino, J.T. 1989. Geographic variation and community structure in an ant-plant mutualism: Azteca and *Cecropia* in Costa Rica. *Biotropica* 21: 126-132.
- McPherson, B.A. & K. Chung-Kim (eds.) 1993. *Evolution of insect pests: patterns of variation*. Wiley, New York.
- Molyneux, R.J., B.C. Campbell, & D.L. Dreyer 1990. Honeydew analysis for detecting phloem transport of plant natural products: implications for host-plant resistance to sap-sucking insects. *Journal of Chemical Ecology* 16: 1899-1909.
- Roth, I. 1976. *Estructura interna de los domacios foliares en Tococa* (Melastomataceae). *Acta Biologica Venezuelana* 9: 221-258.
- Shannon, M.L., T.M. Palmer, & T.P. Young 2002. Competition-colonization trade-offs in a guild of African Acacia-ants. *Ecological Monographs* 72: 347-363.
- Vasconcelos, H.L. & D.W. Davidson 2000. Relationship between plant size and ant associates in two Amazonian ant-plants. *Biotropica* 32: 100-111.
- Way, M.J. 1963. Mutualism between ants and honeydew-producing Homoptera. *Annual Review of Entomology* 8: 307-344.
- Yu, D.W. & D.W. Davidson 1997. Experimental studies of species-specificity in *Cecropia*-ant relationships. *Ecological Monographs* 67: 273-294.

